

Phytoplankton functional types observation from space in the Fram Strait (2002-2020)



Hongyan Xi^{1*}, Ilka Peeken¹, Svetlana Losa¹, Antoine Mangin², Leonardo Alvarado¹, Marine Bretagnon², Astrid Bracher^{1,3}

¹ Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

² ACRI-ST, Sophia Antipolis, France

³ IUP, University of Bremen, Bremen, Germany

* Hongyan.Xi@awi.de

- **Fram Strait - Gateway to the Arctic:** warmer nutrient-rich Atlantic water meets the cold fresher Arctic water
- Complex ecosystem subject to severe climate-induced environmental changes
- Phytoplankton as the base of food web regulate the key biogeochemical processes are highly variable there



Objectives

- Evaluation of satellite PFT products in the Fram Strait
- Time series of PFTs in the Fram Strait from 2002 to 2020 (separating Atlantic and Arctic water masses)
- PFT phenology, inter-annual variations, changes in composition

Evaluation of PFT products from satellites


- A global approach for PFT chlorophyll retrieval using ocean color reflectance data and SST**
 - A set of empirical orthogonal function based algorithms
 - Capability of retrieving Chla of 6 groups- diatoms, haptophytes and others
- Data products available in CMEMS at <https://marine.copernicus.eu/>**

(Xi et al. 2020, 2021)




Matchup of mean of 3x3 pixels (with > 4 valid pixels and CV>0.2) to in-situ

Satellite PFT composites versus in situ PFT during expedition PS74: 20090622-0731 (an example)





PFT observation in the west and east Fram Strait


- Whole region: 16W – 12E, 75N – 81N
- Period: April to August (2002-2020)
- A dynamic clustering of the water masses based on CMEMS daily SST




PFT phenology Bloom maxima date



PFT composition



PFT Climatology



Summary and outlook

- Satellite PFT data match well with in situ pigment derived PFT
- Phytoplankton time series
 - Annual cycles of phyto growth are different in the west and east FS
 - High interannual variability
 - Blooms start slightly earlier in the west FS (more related to the marginal ice zone)
 - Haptophytes grow after diatoms and last until August
 - Obvious changes in the last few years?
- Further investigate the changes found for satellite PGs in linkage to other biogeochemical/physical parameters
- Phytoplankton carbon estimation by accounting for PFT community structure

Acknowledgements

- **ACRI-AWI joint project (OLCI-PFT)**
- EU project PORTWIMS
- EU Copernicus Marine Service (CMEMS)
- Marc Taylor, Sonja Wiegmann and all the AWI PhytoOptics previous and current team members
- PEBCAO team
- All the scientists and crew involved in the global HPLC data collection and analyses for providing their pigment data
- NASA, ESA and EUMETSAT for the SeaWiFS, MODIS, VIIRS, MERIS, and OLCI data, and specially the ACRI-ST GlobColour team for providing the OLCI and merged ocean color L3 products.

Perspective slide

Knowledge gaps (priorities from high to low in a time manner)

- Gaps in satellite PFT products (spatial and temporal) and inconsistency between different sensors (1-2 year)
- Difficulty in validating PFT products (different data type, estimation methods and units) (3-5 years)
- Uncertainty in the in situ pigment-derived PFT (diagnostic pigment analysis, CHEMTAX..) to allow comprehensive uncertainty study of the satellite PFTs (3-5 years)
- Accounting phytoplankton community structure for better phytoplankton carbon estimation and also primary production (3-5 years)
- Well synergy between in situ, satellite and modelling output with respect to phytoplankton community and be able to interpret trends over larger time scales (20 years)